

Subject Code :

V/MAT (v)

Booklet No.

V/MAT (v)

2014

(5th Semester)

MATHEMATICS

Paper : MATH-351

(Computer-oriented Numerical Analysis)

Full Marks : 75

Time : 3 hours

(PART : B—DESCRIPTIVE)

(Marks : 50)

*The figures in the margin indicate full marks
for the questions*

Answer one question from each Unit

UNIT—I

1. (a) Find the negative root of $x^3 - 2x + 5 = 0$ correct to three places of decimals by Newton-Raphson method. 5
- (b) Find the function whose first difference is $x^3 + 3x^2 + 5x + 12$. Write the result in factorial polynomial expansion form. 3+2=5
2. (a) Find the positive real root of $x \log_{10} x = 1.2$ using the bisection method in four iterations. 5

- (b) Find the second difference of the polynomial $f(x) = 7x^4 + 12x^3 - 6x^2 + 5x - 3$ taking $h = 2$ giving the result in normal polynomial form. 4+1=5

UNIT—II

3. (a) Obtain Newton's divided difference interpolation formula for interpolation with non-equal intervals of the argument. 6

- (b) Find $\log_{10} \pi$ by Newton's formula for forward interpolation given that

$$\log_{10} 3.141 = 0.4970679364$$

$$\log_{10} 3.142 = 0.4972061807$$

$$\log_{10} 3.143 = 0.4973443810$$

$$\log_{10} 3.144 = 0.4974825374$$

$$\log_{10} 3.145 = 0.4976206498$$

using $\pi = 3.1415926536$. 4

4. (a) Find the equation of the cubic curve which passes through the points (4, -43), (7, 83), (9, 327), (12, 1053). 4

- (b) Obtain Newton's backward interpolation formula for interpolation with equal intervals of the argument. 6

UNIT—III

5. (a) Solve the following system of equations by Gaussian elimination method : 6

$$2.51x + 1.48y + 4.53z = 0.05$$

$$1.48x + 0.93y - 1.30z = 1.03$$

$$2.68x + 3.04y - 1.48z = -0.53$$

- (b) Solve the following by Gauss-Jordan method : 4

$$x + y + z = 9$$

$$2x - 3y + 4z = 13$$

$$3x + 4y + 5z = 40$$

6. (a) Solve $2x + 3y = 3$; $2x + 3y = 5$ by Gauss-Seidel iteration method. 4

- (b) By Crout's method, solve the following system of simultaneous equations : 6

$$2x - 6y + 8z = 24$$

$$5x + 4y - 3z = 2$$

$$3x + y + 2z = 16$$

UNIT—IV

7. (a) The function $f(x)$ is tabulated in the table below :

x	0.7	0.8	0.9	1.0	1.1
y	0.644218	0.717356	0.783327	0.841471	0.891207

Find $f'(0.77)$.

4

- (b) Obtain the formula for Simpson's one-third rule of integration for numerical integration. 6

8. (a) From the table below, find $f''(0.15)$: 6

x	0.0	0.05	0.1	0.15	0.20	0.25	0.30	0.35
y	1.275	1.342	1.517	1.821	1.99	2.45	2.85	3.40

- (b) Evaluate $\int_0^{10} \frac{dx}{1+x^2}$ by using trapezoidal rule. 4

UNIT—V

9. (a) Given that

$$\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}, \quad y(1) = 1$$

Evaluate $y(1.3)$ by Euler's multiple-step method. 6

- (b) Find the approximate solution of the initial value problem $y' = 1 + y^2$, $y(0) = 0$ by Picard's method and compare it with the exact solution. 4

10. Using any predictor-corrector method, find $y(0.4)$ for the differential equation

$$\frac{dy}{dx} = 1 + xy, \quad y(0) = 2 \quad 10$$

2014

(5th Semester)

MATHEMATICS

Paper : MATH-351

(Computer-oriented Numerical Analysis)

(PART : A—OBJECTIVE)

(Marks : 25)

Answer **all** questions**SECTION—A**

(Multiple Choice)

(Marks : 10)

Each question carries 1 mark

Put a Tick ☒ mark against the correct answer in the box provided :

1. Which of the following statements is true for difference operator, where Δ , ∇ , δ , μ stand for forward, backward, central and average difference operators respectively?

(a) $\Delta^n y_i = \Delta^{n-1} y_{i+1} - \Delta^{n-1} y_i$ ☐

(b) $\nabla^n y_i = \nabla^n y_{i+1} - \nabla^{n-1} y_{i-1}$ ☐

(c) $\delta f(x) = f\left(x - \frac{h}{2}\right) - f\left(x + \frac{h}{2}\right)$ ☐

(d) $\mu f(x) = f\left(x + \frac{h}{2}\right) + f\left(x - \frac{h}{2}\right)$ ☐

(2)

2. A reciprocal factorial function denoted by $x^{(-n)}$, where n is a positive integer is a product of the form given by

(a) $(x+h)(x+2h)(x+3h)\dots(x+nh)$ ☐

(b) $\frac{1}{x(x+h)(x+2h)(x+3h)\dots(x+nh)}$ ☐

(c) $\frac{1}{x(x+h)(x+2h)(x+3h)\dots[x+(n-1)h]}$ ☐

(d) $\frac{1}{(x+h)(x+2h)(x+3h)\dots(x+nh)}$ ☐

3. While constructing a forward difference table if six arguments are given, the forward difference table will contain terms up to

(a) $\Delta^6 y$ ☐

(b) $\Delta^5 y$ ☐

(c) $\Delta^7 y$ ☐

(d) None of the above ☐

4. If $f(x) = \frac{1}{x^2}$, then the divided difference of $\delta(a, b)$ is equal to

(a) $\frac{-(a+b)}{a^2b^2}$ ☐

(b) $\frac{ab+bc+ac}{a^2b^2c^2}$ ☐

(c) $\frac{(a-b)}{a^2b^2}$ ☐

(d) $\frac{(b-a)}{a^2b^2}$ ☐

5. The method for obtaining the solution of the system of simultaneous equation by Gauss elimination method depends on reducing the coefficient matrix to a/an

(a) diagonal matrix ☐

(b) lower triangular matrix ☐

(c) upper triangular matrix ☐

(d) diagonally dominant matrix ☐

6. The coefficient matrix obtained from the simultaneous equations

$$\begin{aligned} a_{11}x + a_{12}y + a_{13}z &= d_1, & b_{21}x + b_{22}y + b_{23}z &= d_2, \\ c_{31}x + c_{32}y + c_{33}z &= d_3 \end{aligned}$$

will be a diagonally dominant matrix if

$$\begin{aligned} &|a_{11}| \geq |a_{12}| + |a_{13}| \\ (a) \quad &|b_{21}| \geq |b_{22}| + |b_{23}| \quad \square \\ &|c_{31}| \geq |c_{32}| + |c_{33}| \end{aligned}$$

$$\begin{aligned} &|a_{11}| \geq |a_{12}| + |a_{13}| \\ (b) \quad &|b_{22}| \geq |b_{21}| + |b_{23}| \quad \square \\ &|c_{33}| \geq |c_{31}| + |c_{32}| \end{aligned}$$

$$\begin{aligned} &|a_{11}| \leq |a_{12}| + |a_{13}| \\ (c) \quad &|b_{22}| \leq |b_{21}| + |b_{23}| \quad \square \\ &|c_{33}| \leq |c_{31}| + |c_{32}| \end{aligned}$$

$$\begin{aligned} &|a_{11}| + |a_{12}| + |a_{13}| \geq |d_1| \\ (d) \quad &|b_{21}| + |b_{22}| + |b_{23}| \geq |d_2| \quad \square \\ &|c_{31}| + |c_{32}| + |c_{33}| \geq |d_3| \end{aligned}$$

7. In the general quadrature formula, trapezoidal rule is obtained by putting

$$(a) \quad n = 2 \quad \square$$

$$(b) \quad n = 4 \quad \square$$

$$(c) \quad n = 2 \text{ and } 4 \text{ both} \quad \square$$

$$(d) \quad n = 1 \quad \square$$

8. The value of $y'(5)$ from the table

x	0	1	2	3	4
y	1	1	15	40	85

is

(a) 625 ☐

(b) 6.25 ☐

(c) 0.625 ☐

(d) 0.0625 ☐

9. The formula for solving the first-order differential equation $\frac{dy}{dx} = f(x, y)$ with $y(x_0) = y_0$ by Taylor's series method is

(a) $y_1 = y_0 + \int_{x_0}^x f(x, y_0) dx$ ☐

(b) $y_1 = y_0 + \frac{h}{1!} y'_0 + \frac{h^2}{2!} y''_0 + \dots$ ☐

(c) $y_1 = y_0 + hf(x_0, y_0)$ ☐

(d) $y_1 = y_0 + \Delta y$ ☐

10. For solving ordinary differential equation numerically, which among the following is applied if successive derivatives can be obtained easily?

- (a) Taylor's method ☐
- (b) Picard's method ☐
- (c) Euler's method ☐
- (d) Runge-Kutta method ☐

(7)

SECTION—B

(Very Short Answer)

(Marks : 15)

Each question carries 3 marks

1. Evaluate

$$\Delta^{10}[(1-x)(1-2x^2)(1-3x^3)(1-4x^4)]$$

where Δ is forward difference operator.

2. Obtain the relation between divided difference and simple difference given by

$$\delta(x_n, x_{n-1}, \dots, x_3, x_2, x_1, x_0) = \frac{\Delta^n y_0}{n! h^n}$$

where δ denotes divided difference and Δ denotes simple difference.

3. What is diagonally dominant system for a simultaneous equation? Are the given equations diagonally dominant?

$$3x + 9y - 2z = 10$$

$$4x + 2y + 13z = 19$$

$$4x - 2y + z = 3$$

4. Use Simpson's 1/3rd rule to find an approximate value of $\log 2$ from the integral $\int_1^2 \frac{dx}{x}$.

5. Using Runge-Kutta method of second-order, find $y(0.1)$ for the differential equation

$$\frac{dy}{dx} = xy + y^2, \quad y(0) = 1$$
